

ON COHOMOLOGICAL DECOMPOSABILITY OF ALMOST-KÄHLER STRUCTURES

DANIELE ANGELLA, ADRIANO TOMASSINI, AND WEIYI ZHANG

ABSTRACT. We study the J -invariant and J -anti-invariant cohomological subgroups of the de Rham cohomology of a compact manifold M endowed with an almost-Kähler structure (J, ω, g) . In particular, almost-Kähler manifolds satisfying a Lefschetz type property, and solvmanifolds endowed with left-invariant almost-complex structures are investigated.

First published in Proceedings of the American Mathematical Society in 2012, published by the American Mathematical Society.

INTRODUCTION

Cohomological properties of compact complex, and, more in general, almost-complex, manifolds have been recently studied by many authors, see, e.g., [3], respectively [11, 12], and the references therein. The study of the cohomology of almost-complex manifolds is motivated, in particular, by a question of Donaldson's, [10, Question 2], relating the tamed and compatible symplectic cones of a compact 4-dimensional almost-complex manifold, see, e.g., [20], and by the analogous question arising for compact higher dimensional complex manifolds, see [20, page 678] and [26, Question 1.7]. (We recall that a symplectic structure ω on a manifold M is said to *tame* an almost-complex structure J if $\omega_x(u_x, J_x u_x) > 0$ for any $x \in M$ and for any $u \in T_x M \setminus \{0\}$, and it is said *compatible* with J if $g := \omega(\cdot, J \cdot)$ is a J -Hermitian metric; in the latter case, the triple (J, ω, g) is called an *almost-Kähler structure* on M .)

Following T.-J. Li and the third author, [20], an almost-complex structure J on a $2n$ -dimensional manifold M is called \mathcal{C}^∞ -*pure-and-full* if

$$H_{dR}^2(M; \mathbb{R}) = H_J^{(1,1)}(M)_{\mathbb{R}} \oplus H_J^{(2,0),(0,2)}(M)_{\mathbb{R}},$$

where $H_J^{(1,1)}(M)_{\mathbb{R}}$ and $H_J^{(2,0),(0,2)}(M)_{\mathbb{R}}$ denote the subgroups of $H_{dR}^2(M; \mathbb{R})$ whose elements can be represented by forms of type $(1, 1)$ and $(2, 0) + (0, 2)$ respectively. In the notation of T. Drăghici, T.-J. Li, and the third author, [11], $H_J^{(1,1)}(M)_{\mathbb{R}} =: H_J^+(M)$ and $H_J^{(2,0),(0,2)}(M)_{\mathbb{R}} =: H_J^-(M)$ are the J -invariant and the J -anti-invariant cohomology subgroups respectively.

In [11, Theorem 2.3], T. Drăghici, T.-J. Li, and the third author proved that every almost-complex structure on a compact 4-dimensional manifold is \mathcal{C}^∞ -pure-and-full. This is no more true in dimension greater than four, see, e.g., [15, Example 3.3], see also [1, 2].

2000 *Mathematics Subject Classification.* 53C55; 53C25; 32G05.

Key words and phrases. \mathcal{C}^∞ -pure-and-full structure; J -anti-invariant cohomology; almost-Kähler structure.

The first and the second authors are partially supported by GNSAGA of INdAM..

The groups $H_J^{(1,1)}(M)_\mathbb{R}$ and $H_J^{(2,0),(0,2)}(M)_\mathbb{R}$ appear as a natural generalization of the Dolbeault cohomology groups to the non-integrable case, see, e.g., [20, Proposition 2.1]. In fact, compact Kähler manifolds are \mathcal{C}^∞ -pure-and-full, and, in this case, $H_J^{(1,1)}(M)_\mathbb{R} \simeq H_{\bar{\partial}}^{1,1}(M) \cap H_{dR}^2(M; \mathbb{R})$ and $H_J^{(2,0),(0,2)}(M)_\mathbb{R} \simeq (H_{\bar{\partial}}^{2,0}(M) \oplus H_{\bar{\partial}}^{0,2}(M)) \cap H_{dR}^2(M; \mathbb{R})$.

We remark that, on a compact complex manifold, other cohomologies can be defined, namely, the Bott-Chern and Aeppli cohomologies. In [3], the problem of cohomology decomposition in terms of the Bott-Chern cohomology groups is investigated, providing in particular a characterization of compact complex manifolds satisfying the $\partial\bar{\partial}$ -Lemma.

Compact Kähler manifolds being \mathcal{C}^∞ -pure-and-full, in this paper we are interested in the study of the cohomological subgroups $H_J^{(1,1)}(M)_\mathbb{R}$ and $H_J^{(2,0),(0,2)}(M)_\mathbb{R}$ for almost-Kähler manifolds.

On the one hand, A. Fino and the second author, [15, Proposition 3.2], as well as T. Drăghici, T.-J. Li, and the third author, [11, Proposition 2.8], proved that the almost-complex structure of a compact almost-Kähler manifold is \mathcal{C}^∞ -pure. On the other hand, we prove the following result, showing therefore a difference between the integrable and the non-integrable cases.

Proposition 4.1. *Let $X := \mathbb{Z}[i]^3 \setminus (\mathbb{C}^3, *)$ be the real manifold underlying the Iwasawa manifold. Then there exists an almost-Kähler structure (J, ω, g) on X which is \mathcal{C}^∞ -pure and non- \mathcal{C}^∞ -full. Furthermore, the Lefschetz type operator $\mathcal{L}_\omega := \omega \wedge \cdot : \wedge^2 M \rightarrow \wedge^4 M$ of the almost-Kähler structure (J, ω, g) does not take g -harmonic 2-forms to g -harmonic 4-forms.*

In studying cohomological decomposition of the de Rham cohomology of almost-Kähler manifolds, the third author introduced a *Lefschetz type property* for 2-forms, see Definition 2.2. Such a property is stronger than the Hard Lefschetz Condition on 2-classes, namely, the property that $[\omega]^{n-2} \smile \cdot : H_{dR}^2(M; \mathbb{R}) \rightarrow H_{dR}^{2n-2}(M; \mathbb{R})$ is an isomorphism, where $2n := \dim M$.

We study such a Lefschetz type property on almost-Kähler manifolds (M, J, ω, g) in relation to the existence of a cohomological decomposition of $H_{dR}^2(M; \mathbb{R})$. More precisely, we prove the following result.

Theorem 2.3. *Let (M, J, ω, g) be a compact almost-Kähler manifold. Suppose that there exists a basis of $H_{dR}^2(X; \mathbb{R})$ represented by g -harmonic 2-forms which are of pure type with respect to J . Then the Lefschetz type property on 2-forms is satisfied.*

Note that, by the hypothesis, it follows, in particular, that J is \mathcal{C}^∞ -pure-and-full and pure-and-full, [15, Theorem 3.7]. Note also that A. Fino and the second author provided in [15] several examples of compact non-Kähler solvmanifolds admitting a basis of harmonic representatives of pure-type with respect to the almost-complex structure. In [13, §2], T. Drăghici, T.-J. Li, and the third author ask whether such a Lefschetz type property on 2-forms is actually equivalent to \mathcal{C}^∞ -fullness for every almost-Kähler nilmanifold and solvmanifold, without any further assumption; Theorem 2.3 and Proposition 4.1 provide results and examples in favour of a possibly positive answer to their question.

In [12, Theorem 1.1], starting with a compact complex surface (M, J) , it is shown that the dimension $h_{\bar{J}}^-$ of the \bar{J} -anti-invariant cohomology subgroup $H_{\bar{J}}^-(M)$

of any *metric related* almost-complex structure \tilde{J} on M (namely, an almost-complex structure \tilde{J} on M inducing the same orientation as that one induced by J and with a common compatible metric), such that $\tilde{J} \neq \pm J$, can be 0, 1, or 2, and a description of such almost-complex structures \tilde{J} having $h_{\tilde{J}}^- \in \{1, 2\}$ is provided. Furthermore, it is conjectured that $h_{\tilde{J}}^- = 0$ for a generic almost-complex structure J on a compact 4-dimensional manifold, and that if $h_{\tilde{J}}^- \geq 3$, then J is integrable, [12, Conjecture 2.4, Conjecture 2.5]. One could set a similar question for higher dimensional manifolds, asking Question 5.2: *are there examples of non-integrable almost-complex structures J on a compact $2n$ -dimensional manifold with $h_{\tilde{J}}^- > n(n-1)$?*

Finally, we prove a Nomizu-type result for the subgroups $H_J^\pm(M)$ of a completely-solvable solvmanifolds $M = \Gamma \backslash G$ endowed with left-invariant almost-complex structures J . More precisely, denote the Lie algebra associated to G by \mathfrak{g} , and consider

$$H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}} := \left\{ \mathfrak{a} = [\alpha] \in H^\bullet(\wedge^\bullet \mathfrak{g}^*, d) : \alpha \in \wedge_J^{(p,q),(q,p)} \mathfrak{g}^* \right\} \subseteq H_{dR}^\bullet(M; \mathbb{R})$$

the subgroup of $H_{dR}^\bullet(M; \mathbb{R})$ that consists of classes admitting a left-invariant representative of type $(p, q) + (q, p)$, where $\wedge_J^{(p,q),(q,p)} \mathfrak{g}^* := (\wedge^{p,q}(\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C})^* \oplus \wedge^{q,p}(\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C})^*) \cap \wedge^\bullet \mathfrak{g}^*$; then the following result holds.

Theorem 5.4. *Let $M = \Gamma \backslash G$ be a solvmanifold endowed with a left-invariant almost-complex structure J , and denote the Lie algebra naturally associated to G by \mathfrak{g} . For any $p, q \in \mathbb{N}$, the map $j: H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}} \rightarrow H_J^{(p,q),(q,p)}(M)_{\mathbb{R}}$ induced by left-translations is injective, and, if $H_{dR}^\bullet(\wedge^\bullet \mathfrak{g}^*, d) \simeq H_{dR}^\bullet(M; \mathbb{R})$ (for instance, if M is a completely-solvable solvmanifold), then $j: H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}} \rightarrow H_J^{(p,q),(q,p)}(M)_{\mathbb{R}}$ is in fact an isomorphism.*

In particular, it follows that $\dim_{\mathbb{R}} H_J^-(M) \leq n(n-1)$ for every left-invariant almost-complex structure on a completely-solvable solvmanifold.

Acknowledgements. The authors would like to thank Tedi Drăghici for useful comments and remarks and Tian-Jun Li for helpful discussions. The first author would like to thank also Jean-Pierre Demailly for useful conversations and for his warm hospitality at Institut Fourier, Université de Grenoble I. The second author would like to thank the Department of Mathematics of University of Notre Dame and the School of Mathematics of University of Minnesota for their warm hospitality. The third author would like to thank Institut des Hautes Études Scientifiques for providing excellent research environment. We are also pleased to thank the referee for fruitful suggestions and remarks.

1. \mathcal{C}^∞ -PURE-AND-FULL ALMOST-COMPLEX STRUCTURES

1.1. Subgroups of the de Rham cohomology of an almost-complex manifold. We start by fixing some notation and recalling some recent results on cohomological properties of almost-complex manifolds; for more details see, e.g., [20, 11, 12, 15, 1, 2, 13], and the references therein.

Let J be a smooth almost-complex structure on a compact $2n$ -dimensional manifold M . Denote by $\wedge^r M$ the bundle of r -forms on M ; we denote with the same symbol $\wedge^r M := \Gamma(M, \wedge^r M)$ the space of smooth global sections of the bundle $\wedge^r M$. Then J extends to a complex automorphism of $T^{\mathbb{C}}M = TM \otimes \mathbb{C}$ such that $T^{\mathbb{C}}M = T_J^{1,0}M \oplus T_J^{0,1}M$, where $T_J^{1,0}M$ and $T_J^{0,1}M$ are the $(\pm i)$ -eigenbundles. The action of J can be extended to the space $\wedge^r(M; \mathbb{C})$ of smooth global sections of the

bundle $\wedge^r(M; \mathbb{C}) := \wedge^r M \otimes \mathbb{C}$ getting the following decomposition:

$$\wedge^r(M; \mathbb{C}) = \bigoplus_{p+q=r} \wedge_J^{p,q} M.$$

Then the space $\wedge^r M$ of real smooth differential r -forms decomposes as

$$\wedge^r M = \bigoplus_{p+q=r, p \leq q} \wedge_J^{(p,q),(q,p)}(M)_{\mathbb{R}},$$

where, for $p < q$, (later on, we do not distinguish the cases $p < q$ and $p = q$)

$$\wedge_J^{(p,q),(q,p)}(M)_{\mathbb{R}} := \{\alpha \in \wedge_J^{p,q} M \oplus \wedge_J^{q,p} M : \alpha = \overline{\alpha}\}, \quad \wedge_J^{(p,p)}(M)_{\mathbb{R}} := \{\alpha \in \wedge_J^{p,p} M : \alpha = \overline{\alpha}\}$$

In particular, for $r = 2$, we will adopt the following notation:

$$\wedge_J^{1,1}(M)_{\mathbb{R}} =: \wedge_J^+ M, \quad \wedge_J^{(2,0),(0,2)}(M)_{\mathbb{R}} =: \wedge_J^- M;$$

this is consistent with the decomposition in invariant and anti-invariant part of $\wedge^2 M$ under the natural action of J on $\wedge^2 M$, given by $J\alpha(\cdot, \cdot) := \alpha(J\cdot, J\cdot)$.

We will refer to forms in $\wedge_J^{1,1}(M)_{\mathbb{R}}$, respectively $\wedge_J^{(2,0),(0,2)}(M)_{\mathbb{R}}$ as forms of *pure type with respect to J* .

For a finite set S of pairs of integers, let

$$\mathcal{Z}_J^S := \bigoplus_{(p,q) \in S, p \leq q} \mathcal{Z}_J^{(p,q),(q,p)}, \quad \mathcal{B}_J^S := \bigoplus_{(p,q) \in S, p \leq q} \mathcal{B}_J^{(p,q),(q,p)},$$

where

$$\mathcal{Z}_J^{(p,q),(q,p)} := \left\{ \alpha \in \wedge_J^{(p,q),(q,p)}(M)_{\mathbb{R}} : d\alpha = 0 \right\},$$

$$\mathcal{B}_J^{(p,q),(q,p)} := \left\{ \beta \in \wedge_J^{(p,q),(q,p)}(M)_{\mathbb{R}} : \text{there exists } \gamma \text{ such that } d\gamma = \beta \right\}.$$

Define

$$H_J^S(M)_{\mathbb{R}} := \frac{\mathcal{Z}_J^S}{\mathcal{B}_J^S}.$$

Let \mathcal{B} be the space of d-exact forms. Since $\frac{\mathcal{Z}_J^S}{\mathcal{B}_J^S} = \frac{\mathcal{Z}_J^S}{\mathcal{B} \cap \mathcal{Z}_J^S}$, a natural inclusion $\rho_S: \frac{\mathcal{Z}_J^S}{\mathcal{B}_J^S} \rightarrow \frac{\mathcal{Z}_J^S}{\mathcal{B}}$ is defined. As in [20], we will write $\rho_S \left(\frac{\mathcal{Z}_J^S}{\mathcal{B}_J^S} \right)$ simply as $\frac{\mathcal{Z}_J^S}{\mathcal{B}}$ and consequently the cohomology spaces $H_J^S(M)_{\mathbb{R}}$ can be identified as

$$H_J^S(M)_{\mathbb{R}} = \{[\alpha] \in H_{dR}^\bullet(M; \mathbb{R}) : \alpha \in \mathcal{Z}_J^S\} = \frac{\mathcal{Z}_J^S}{\mathcal{B}}.$$

Therefore, there is a natural inclusion

$$H_J^{(1,1)}(M)_{\mathbb{R}} + H_J^{(2,0),(0,2)}(M)_{\mathbb{R}} \subseteq H_{dR}^2(M; \mathbb{R}).$$

1.2. \mathcal{C}^∞ -pure-and-full and pure-and-full almost-complex structures. As in [20], we set the following definition.

Definition 1.1 ([20, Definition 2.2, Definition 2.3, Lemma 2.2]). An almost-complex structure J on a manifold M is said to be

- \mathcal{C}^∞ -pure if $H_J^{(1,1)}(M)_{\mathbb{R}} \cap H_J^{(2,0),(0,2)}(M)_{\mathbb{R}} = \{0\}$,
- \mathcal{C}^∞ -full if $H_{dR}^2(M; \mathbb{R}) = H_J^{(1,1)}(M)_{\mathbb{R}} + H_J^{(2,0),(0,2)}(M)_{\mathbb{R}}$,
- \mathcal{C}^∞ -pure-and-full if

$$H_{dR}^2(M; \mathbb{R}) = H_J^{(1,1)}(M)_{\mathbb{R}} \oplus H_J^{(2,0),(0,2)}(M)_{\mathbb{R}}.$$

According to the previous notation, we will write

$$H_J^+(M) := H_J^{(1,1)}(M)_{\mathbb{R}}, \quad H_J^-(M) := H_J^{(2,0),(0,2)}(M)_{\mathbb{R}}.$$

Similar definitions in terms of currents can be given, introducing the notion of *pure-and-full* almost-complex structure: we refer to [20, §2.2.2] for further details and results. More precisely, on an almost complex manifold (M, J) , the space $\mathcal{E}_k(M)_{\mathbb{R}}$ of real k -currents has a decomposition $\mathcal{E}_k(M)_{\mathbb{R}} = \bigoplus_{p+q=k} \mathcal{E}_{(p,q),(q,p)}^J(M)_{\mathbb{R}}$, where $\mathcal{E}_{(p,q),(q,p)}^J(M)_{\mathbb{R}}$ denotes the space of real k -currents of bi-dimension $(p, q) + (q, p)$.

Let $\mathcal{Z}_{(2,0),(0,2)}^J$ and $\mathcal{Z}_{(1,1)}^J$ denote the spaces of real d-closed currents of bi-dimension $(2, 0) + (0, 2)$, respectively $(1, 1)$, and $\mathcal{B}_{(2,0),(0,2)}^J$ and $\mathcal{B}_{(1,1)}^J$ denote the spaces of real d-exact currents of bi-dimension $(2, 0) + (0, 2)$, respectively $(1, 1)$. Denote by \mathcal{B} the space of boundaries. Let, as in [20],

$$H_{(1,1)}^J(M)_{\mathbb{R}} := \left\{ [\alpha] \in H_2(M; \mathbb{R}) : \alpha \in \mathcal{Z}_{(1,1)}^J \right\} = \frac{\mathcal{Z}_{(1,1)}^J}{\mathcal{B}},$$

$$H_{(2,0),(0,2)}^J(M)_{\mathbb{R}} := \left\{ [\alpha] \in H_2(M; \mathbb{R}) : \alpha \in \mathcal{Z}_{(2,0),(0,2)}^J \right\} = \frac{\mathcal{Z}_{(2,0),(0,2)}^J}{\mathcal{B}}.$$

We recall the following definition.

Definition 1.2 ([20, Definition 2.15, Definition 2.16]). An almost complex structure J on a manifold M is said to be *pure* if $H_{(1,1)}^J(M)_{\mathbb{R}} \cap H_{(2,0),(0,2)}^J(M)_{\mathbb{R}} = \{0\}$. It is said to be *full* if $H_2(M; \mathbb{R}) = H_{(1,1)}^J(M)_{\mathbb{R}} + H_{(2,0),(0,2)}^J(M)_{\mathbb{R}}$. Therefore, an almost complex structure J is *pure-and-full* if and only if

$$H_2(M, \mathbb{R}) = H_{(1,1)}^J(M)_{\mathbb{R}} \oplus H_{(2,0),(0,2)}^J(M)_{\mathbb{R}}.$$

In [20, Proposition 2.1] it is shown that, given a compact complex manifold (M, J) of complex dimension n , if $n = 2$ or J is Kähler, then J is \mathcal{C}^∞ -pure-and-full, and $H_J^{(1,1)}(M)_{\mathbb{R}} \simeq H_{\bar{\partial}}^{1,1}(M) \cap H_{dR}^2(M; \mathbb{R})$ and $H_J^{(2,0),(0,2)}(M)_{\mathbb{R}} \simeq \left(H_{\bar{\partial}}^{2,0}(M) \oplus H_{\bar{\partial}}^{0,2}(M) \right) \cap H_{dR}^2(M; \mathbb{R})$. In view of this result, the subgroups $H_J^{(1,1)}(M)_{\mathbb{R}}$ and $H_J^{(2,0),(0,2)}(M)_{\mathbb{R}}$ of the de Rham cohomology can be viewed as an analogue of the Dolbeault cohomology groups for non-integrable almost-complex structures.

In [11, Theorem 2.3] it is proven the following result.

Theorem 1.3 ([11, Theorem 2.3]). *If M is a compact manifold of dimension 4, then any almost-complex structure J on M is \mathcal{C}^∞ -pure-and-full.*

This is no more true in dimension higher than 4: in [15, Example 3.3], a compact non- \mathcal{C}^∞ -pure almost-complex structure on a 6-dimensional nilmanifold is constructed. Therefore, the previous theorem can be considered a sort of Hodge decomposition theorem in the non-Kähler case.

2. COHOMOLOGICAL PROPERTIES OF ALMOST-KÄHLER MANIFOLDS

2.1. Lefschetz type property on almost-Kähler manifolds with pure-type harmonic representatives. Given a compact $2n$ -dimensional almost-Kähler manifold (M, J, ω, g) , we are interested in studying the property of being \mathcal{C}^∞ -pure-and-full.

Firstly we recall the following result.

Proposition 2.1 ([11, Proposition 2.8], [15, Proposition 3.2]). *If J is an almost-complex structure on a compact manifold M and J admits a compatible symplectic structure, then J is \mathcal{C}^∞ -pure.*

Furthermore, A. Fino and the second author proved that an almost-Kähler manifold admitting a basis of harmonic 2-forms whose elements are of pure type with respect to the almost-complex structure is \mathcal{C}^∞ -pure-and-full and pure-and-full, [15, Theorem 3.7]; they also provided several examples of compact non-Kähler solvmanifolds satisfying the above assumption in [15].

To the purpose of studying the property of being \mathcal{C}^∞ -pure-and-full on almost-Kähler manifolds, we recall the following definition.

Definition 2.2. Given a compact $2n$ -dimensional symplectic manifold (M, ω) , denote by

$$\mathcal{L}_\omega: \wedge^2 M \rightarrow \wedge^{2n-2} M, \quad \mathcal{L}_\omega(\alpha) := \omega^{n-2} \wedge \alpha,$$

the *Lefschetz type operator* (on 2-forms) associated with ω .

Then one says that the compact $2n$ -dimensional almost-Kähler manifold (M, J, ω, g) satisfies the *Lefschetz type property (on 2-forms)* if \mathcal{L}_ω takes g -harmonic 2-forms to g -harmonic $(2n-2)$ -forms.

Furthermore, we recall some notions and results from [6, 22, 27], see also [23, 7]. Let (M, ω) be a compact $2n$ -dimensional symplectic manifold. Extend $\omega^{-1}: T^*M \rightarrow TM$ to the whole exterior algebra of T^*M . For any $k \in \mathbb{N}$, the *symplectic \star_ω operator* is defined as

$$\star_\omega: \wedge^k M \rightarrow \wedge^{2n-k} M, \quad \beta \wedge \star_\omega \alpha = \omega^{-1}(\alpha, \beta) \frac{\omega^n}{n!}, \quad \forall \alpha, \beta \in \wedge^k M.$$

One can prove that $\star_\omega^2 = \text{id}_{\wedge^\bullet M}$, [6, Lemma 2.1.2].

For any $k \in \mathbb{N}$, define the *symplectic co-differential operator*

$$\delta_\omega: \wedge^k M \rightarrow \wedge^{k-1} M, \quad \delta_\omega|_{\wedge^k M} := (-1)^{k+1} \star_\omega d \star_\omega;$$

this operator has been studied by J.-L. Brylinski in [6] for Poisson manifolds; in the context of generalized complex geometry, see, e.g., [16], it can be interpreted as the symplectic counterpart of the operator $d^c := -i(\partial - \bar{\partial})$ in complex geometry, see [7].

By definition, (M, ω) satisfies the *Hard Lefschetz Condition* if, for each $k \in \mathbb{N}$, the map

$$[\omega]^k \smile \cdot: H_{dR}^{n-k}(M; \mathbb{R}) \rightarrow H_{dR}^{n+k}(M; \mathbb{R})$$

is an isomorphism. O. Mathieu, [22, Corollary 2], and, independently, D. Yan, [27, Theorem 0.1], proved that, given a compact symplectic manifold (M, ω) , any de Rham cohomology class has a (possibly non-unique) ω -*symplectically harmonic representative* (that is, a d -closed δ_ω -closed representative) if and only if the Hard Lefschetz Condition holds.

We can now prove the following result.

Theorem 2.3. *Let (M, J, ω, g) be a compact almost-Kähler manifold. Suppose that there exists a basis of $H_{dR}^2(X; \mathbb{R})$ represented by g -harmonic 2-forms which are of pure type with respect to J . Then the Lefschetz type property on 2-forms is satisfied.*

Proof. Recall that, on a $2n$ -dimensional almost-Kähler manifold (M, J, ω, g) , the Hodge $*_g$ operator and the symplectic \star_ω operator are related by $\star_\omega = *_g J$, [6, Theorem 2.4.1, Remark 2.4.4]. Therefore, for forms of pure type with respect to J , the properties of being g -harmonic and of being ω -symplectically harmonic are equivalent. The theorem follows noting that, [27, Lemma 1.2], $[\mathcal{L}_\omega, d] = 0$ and $[\mathcal{L}_\omega, \delta_\omega] = d$, hence \mathcal{L}_ω sends ω -symplectically harmonic 2-forms (of pure type with respect to J) to ω -symplectically harmonic $(2n-2)$ -forms (of pure type with respect to J). \square

Remark 2.4. We note that if (M, J, ω, g) is a compact $2n$ -dimensional almost-Kähler manifold satisfying the Lefschetz type property on 2-forms and J is \mathcal{C}^∞ -full, then J is \mathcal{C}^∞ -pure-and-full and pure-and-full.

Indeed, we have already remarked that J is \mathcal{C}^∞ -pure, see Proposition 2.1. Moreover, since J is \mathcal{C}^∞ -full, J is also pure by [20, Proposition 2.5]. We recall now the argument in [15] to prove that J is also full.

Firstly, note that if the Lefschetz type property on 2-forms holds, then $[\omega^{n-2}] \smile \cdot : H_{dR}^{2n-2}(M; \mathbb{R}) \rightarrow H_{dR}^{2n-2}(M; \mathbb{R})$ is an isomorphism. Therefore, we get that

$$H_{dR}^{2n-2}(M; \mathbb{R}) = H_J^{(n,n-2),(n-2,n)}(M)_{\mathbb{R}} + H_J^{(n-1,n-1)}(M)_{\mathbb{R}} ;$$

indeed, (following the argument in [15, Theorem 4.1],) since $[\omega^{n-2}] \smile \cdot : H_{dR}^{2n-2}(M; \mathbb{R}) \rightarrow H_{dR}^{2n-2}(M; \mathbb{R})$ is in particular surjective, we have

$$\begin{aligned} H_{dR}^{2n-2}(M; \mathbb{R}) &= [\omega^{n-2}] \smile H_{dR}^{2n-2}(M; \mathbb{R}) = [\omega^{n-2}] \smile \left(H_J^{(2,0),(0,2)}(M)_{\mathbb{R}} \oplus H_J^{(1,1)}(M)_{\mathbb{R}} \right) \\ &\subseteq H_J^{(n,n-2),(n-2,n)}(M)_{\mathbb{R}} + H_J^{(n-1,n-1)}(M)_{\mathbb{R}} , \end{aligned}$$

yielding the above decomposition of $H_{dR}^{2n-2}(M; \mathbb{R})$. Then, it follows that J is also full, see, for example, [1, Theorem 2.1].

2.2. A family of almost-Kähler manifolds satisfying the Lefschetz type property on 2-forms. Let \mathfrak{n} be the 6-dimensional nilpotent Lie algebra whose structure equations, with respect to a basis $\{e^j\}_{j \in \{1, \dots, 6\}}$ of \mathfrak{n}^* , are given by

$$de^1 = de^2 = de^3 = 0, \quad de^4 = e^{23}, \quad de^5 = e^{13}, \quad de^6 = e^{12}$$

(where we write e^{jk} instead of $e^j \wedge e^k$). Using a result by Mal'tsev, [21, Theorem 7], the connected simply-connected Lie group G associated with \mathfrak{n} admits a discrete co-compact subgroup Γ : let $N := \Gamma \backslash G$ be the (compact) nilmanifold obtained as a quotient of G by Γ . Note that N is not formal by a theorem of K. Hasegawa's, [17, Theorem 1, Corollary].

Fix $\alpha > 1$ and take

$$\omega_\alpha := e^{14} + \alpha \cdot e^{25} + (\alpha - 1) \cdot e^{36} ;$$

since $d\omega_\alpha = 0$ and $\omega_\alpha^3 \neq 0$, we get that ω_α is a left-invariant symplectic form on N . Set

$$\begin{aligned} J_\alpha e_1 &:= e_4, & J_\alpha e_2 &:= \alpha e_5, & J_\alpha e_3 &:= (\alpha - 1) e_6, \\ J_\alpha e_4 &:= -e_1, & J_\alpha e_5 &:= -\frac{1}{\alpha} e_2, & J_\alpha e_6 &:= -\frac{1}{\alpha-1} e_3, \end{aligned}$$

where $\{e_1, \dots, e_6\}$ denotes the global dual frame of $\{e^1, \dots, e^6\}$ on N . It is immediate to check that

- setting $g_\alpha(\cdot, \cdot) := \omega_\alpha(\cdot, J_\alpha \cdot)$, the triple $(J_\alpha, \omega_\alpha, g_\alpha)$ gives rise to a family of left-invariant almost-Kähler structures on N ;
- denoting by

$$\begin{aligned} E_\alpha^1 &:= e^1, & E_\alpha^2 &:= \alpha e^2, & E_\alpha^3 &:= (\alpha - 1) e^3, \\ E_\alpha^4 &:= e^4, & E_\alpha^5 &:= e^5, & E_\alpha^6 &:= e^6, \end{aligned}$$

then $\{E_\alpha^1, \dots, E_\alpha^6\}$ is a g_α -orthonormal co-frame on N ; with respect to this new co-frame, we easily obtain the following structure equations:

$$d E_\alpha^1 = d E_\alpha^2 = d E_\alpha^3 = 0, \quad d E_\alpha^4 = \frac{1}{\alpha(\alpha - 1)} E_\alpha^{23}, \quad d E_\alpha^5 = \frac{1}{\alpha - 1} E_\alpha^{13}, \quad d E_\alpha^6 = \frac{1}{\alpha} E_\alpha^{12}.$$

Then,

$$\varphi_\alpha^1 := E_\alpha^1 + i E_\alpha^4, \quad \varphi_\alpha^2 := E_\alpha^2 + i E_\alpha^5, \quad \varphi_\alpha^3 := E_\alpha^3 + i E_\alpha^6,$$

are $(1, 0)$ -forms with respect to the almost-complex structure J_α , and

$$\omega_\alpha = E_\alpha^{14} + E_\alpha^{25} + E_\alpha^{36}.$$

By a result of K. Nomizu's, [25, Theorem 1], see Theorem 5.3, the de Rham cohomology of N is straightforwardly computed:

$$H_{dR}^2(N; \mathbb{R}) \simeq \mathbb{R} \left\langle E_\alpha^{15}, E_\alpha^{16}, E_\alpha^{24}, E_\alpha^{26}, E_\alpha^{34}, E_\alpha^{35}, E_\alpha^{14} + \frac{1}{\alpha} E_\alpha^{25}, \frac{1}{\alpha} E_\alpha^{25} + \frac{1}{\alpha - 1} E_\alpha^{36} \right\rangle$$

(where we have listed the g_α -harmonic representatives instead of their classes).

Note that the listed g_α -harmonic representatives of $H_{dR}^2(N; \mathbb{R})$ are of pure type with respect to J_α : hence, the almost-complex structure J_α is \mathcal{C}^∞ -pure-and-full by [15, Theorem 3.7]; in particular, note that

$$\begin{aligned} H_{dR}^2(N; \mathbb{R}) \simeq \mathbb{R} \left\langle i \alpha \varphi_\alpha^{1\bar{1}} + i \varphi_\alpha^{2\bar{2}}, i(\alpha - 1) \varphi_\alpha^{2\bar{2}} + i \alpha \varphi_\alpha^{3\bar{3}}, \Im \varphi_\alpha^{1\bar{2}}, \Im \varphi_\alpha^{1\bar{3}}, \Im \varphi_\alpha^{3\bar{2}} \right\rangle \\ \oplus \langle \Im \varphi_\alpha^{1\bar{2}}, \Im \varphi_\alpha^{1\bar{3}}, \Im \varphi_\alpha^{2\bar{3}} \rangle, \end{aligned}$$

hence $h_{J_\alpha}^+(N) = 5$ and $h_{J_\alpha}^-(N) = 3$.

Moreover, one explicitly notes that

$$\begin{aligned} \mathcal{L}_{\omega_\alpha} E_\alpha^{15} &= E_\alpha^{1536} = *_{g_\alpha} E_\alpha^{24}, & \mathcal{L}_{\omega_\alpha} E_\alpha^{16} &= E_\alpha^{1625} = *_{g_\alpha} E_\alpha^{34}, \\ \mathcal{L}_{\omega_\alpha} E_\alpha^{24} &= E_\alpha^{2436} = *_{g_\alpha} E_\alpha^{15}, & \mathcal{L}_{\omega_\alpha} E_\alpha^{26} &= E_\alpha^{2614} = *_{g_\alpha} E_\alpha^{35}, \\ \mathcal{L}_{\omega_\alpha} E_\alpha^{34} &= E_\alpha^{3425} = *_{g_\alpha} E_\alpha^{16}, & \mathcal{L}_{\omega_\alpha} E_\alpha^{35} &= E_\alpha^{3514} = *_{g_\alpha} E_\alpha^{26}, \end{aligned}$$

while

$$\mathcal{L}_{\omega_\alpha} \left(E_\alpha^{14} + \frac{1}{\alpha} E_\alpha^{25} \right) = -\frac{\alpha + 1}{\alpha} E_\alpha^{1245} - \frac{1}{\alpha} E_\alpha^{2356} - E_\alpha^{1346}$$

where

$$d *_{g_\alpha} \mathcal{L}_{\omega_\alpha} \left(E_\alpha^{14} + \frac{1}{\alpha} E_\alpha^{25} \right) = d \left(-\frac{\alpha + 1}{\alpha} E_\alpha^{36} - E_\alpha^{25} - \frac{1}{\alpha} E_\alpha^{14} \right) = 0,$$

and, by a similar computation, $d *_{g_\alpha} \mathcal{L}_{\omega_\alpha} (e^{25} + e^{36}) = 0$. This proves explicitly that ω_α satisfies the Lefschetz type property on 2-forms.

The nilmanifold N is not formal by a theorem of K. Hasegawa's, [17, Theorem 1, Corollary]. The non-formality of M can be also proved by giving a non-zero triple Massey product on N , see [9]: since

$$[E_\alpha^1] \smile [E_\alpha^3] = (\alpha - 1) [d E_\alpha^5] = 0, \quad [E_\alpha^3] \smile [E_\alpha^2] = -\alpha (\alpha - 1) [d E_\alpha^4] = 0,$$

we get that the triple Massey product

$$\langle [E_\alpha^1], [E_\alpha^3], [E_\alpha^2] \rangle = -(\alpha - 1) [E_\alpha^{25} + \alpha E_\alpha^{14}]$$

does not vanish, and hence N is not formal.

In summary, we have proven the following result.

Proposition 2.5. *There is a non-formal 6-dimensional nilmanifold N endowed with a 1-parameter family $\{(J_\alpha, \omega_\alpha, g_\alpha)\}_{\alpha > 1}$ of left-invariant almost-Kähler structures being C^∞ -pure-and-full and pure-and-full and satisfying the Lefschetz type property on 2-forms.*

Remark 2.6. It has to be noted that $\omega_\alpha \wedge \cdot : \wedge^2 N^6 \rightarrow \wedge^4 N^6$ induces an isomorphism in cohomology $[\omega_\alpha] \smile \cdot : H_{dR}^2(N, \mathbb{R}) \rightarrow H_{dR}^4(N, \mathbb{R})$, while, accordingly to [5, Theorem A], $[\omega_\alpha]^2 \smile \cdot : H_{dR}^1(N, \mathbb{R}) \rightarrow H_{dR}^5(N, \mathbb{R})$ is not an isomorphism.

3. ALMOST-KÄHLER C^∞ -PURE-AND-FULL STRUCTURES

3.1. The Nakamura manifold of completely solvable type. Take $A \in \mathrm{SL}(2; \mathbb{Z})$ with two different real eigenvalues e^λ and $e^{-\lambda}$ with $\lambda > 0$, and fix $P \in \mathrm{GL}(2; \mathbb{R})$ such that $PAP^{-1} = \mathrm{diag}(e^\lambda, e^{-\lambda})$. For example, take

$$A := \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \quad \text{and} \quad P := \begin{pmatrix} \frac{1-\sqrt{5}}{2} & 1 \\ 1 & \frac{\sqrt{5}-1}{2} \end{pmatrix}$$

and consequently $\lambda = \log \frac{3+\sqrt{5}}{2}$. Let $M^6 := M^6(\lambda)$ be the compact complex manifold

$$M^6 := \mathbb{S}_{x^2}^1 \times \frac{\mathbb{R}_{x^1} \times \mathbb{T}_{\mathbb{C}, (x^3, x^4, x^5, x^6)}^2}{\langle T_1 \rangle}$$

where $\mathbb{T}_{\mathbb{C}}^2$ is the 2-dimensional complex torus $\mathbb{T}_{\mathbb{C}}^2 := \frac{\mathbb{C}^2}{P\mathbb{Z}[i]^2}$ and T_1 acts on $\mathbb{R} \times \mathbb{T}_{\mathbb{C}}^2$ as $T_1(x^1, x^3, x^4, x^5, x^6) := (x^1 + \lambda, e^{-\lambda} x^3, e^\lambda x^4, e^{-\lambda} x^5, e^\lambda x^6)$. The manifold M^6 can be seen as a compact quotient of a completely-solvable Lie group by a discrete co-compact subgroup, [14, Example 3.1]; (denote the Lie algebra naturally associated to the completely-solvable Lie group of M^6 by \mathfrak{g}). Using coordinates x^2 on \mathbb{S}^1 , x^1 on \mathbb{R} and (x^3, x^4, x^5, x^6) on $\mathbb{T}_{\mathbb{C}}^2$, we set

$$e^1 := dx^1, \quad e^2 := dx^2, \quad e^3 := e^{x^1} dx^3, \quad e^4 := e^{-x^1} dx^4, \quad e^5 := e^{x^1} dx^5, \quad e^6 := e^{-x^1} dx^6$$

as a basis for \mathfrak{g}^* ; therefore, with respect to $\{e^i\}_{i \in \{1, \dots, 6\}}$, the structure equations are the following:

$$de^1 = de^2 = 0, \quad de^3 = e^{13}, \quad de^4 = -e^{14}, \quad de^5 = e^{15}, \quad de^6 = -e^{16}.$$

3.2. The de Rham cohomology of the Nakamura manifold. Let J be the almost-complex structure on M^6 defined by the complex $(1, 0)$ -forms given by

$$\varphi^1 := \frac{1}{2}(e^1 + ie^2), \quad \varphi^2 := e^3 + ie^5, \quad \varphi^3 := e^4 + ie^6.$$

It is straightforward to check that J is integrable.

Being M^6 a compact quotient of a completely-solvable Lie group, one computes the

de Rham cohomology of M^6 easily by A. Hattori's theorem [19, Corollary 4.2], see Theorem 5.3:

$$\begin{aligned} H_{dR}^1(M^6; \mathbb{C}) &\simeq \mathbb{C} \langle \varphi^1, \bar{\varphi}^1 \rangle, & H_{dR}^2(M^6; \mathbb{C}) &\simeq \mathbb{C} \langle \varphi^{1\bar{1}}, \varphi^{2\bar{3}}, \varphi^{3\bar{2}}, \varphi^{23}, \varphi^{\bar{2}\bar{3}} \rangle, \\ H_{dR}^3(M^6; \mathbb{C}) &\simeq \mathbb{C} \langle \varphi^{12\bar{3}}, \varphi^{13\bar{2}}, \varphi^{123}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{3\bar{1}\bar{2}}, \varphi^{23\bar{1}}, \varphi^{\bar{1}\bar{2}\bar{3}} \rangle \end{aligned}$$

(for the sake of clearness, we write, for example, $\varphi^{A\bar{B}}$ in place of $\varphi^A \wedge \bar{\varphi}^B$ and we list the harmonic representatives with respect to the metric $g := \sum_{j=1}^3 \varphi^j \odot \bar{\varphi}^j$ instead of their classes). Therefore, M^6 is *geometrically formal*, i.e., the product of g -harmonic forms is still g -harmonic, and therefore it is *formal*, namely the de Rham complex of M is formal as a differential graded algebra, see, e.g., [9]. Furthermore, it can be easily checked that

$$\omega := e^{12} + e^{34} + e^{56}$$

gives rise to a symplectic structure on M^6 satisfying the Hard Lefschetz Condition. We obtain the following result.

Proposition 3.1 ([14, Proposition 3.2]). *The manifold M^6 is formal and it admits a symplectic form ω satisfying the Hard Lefschetz Condition.*

Note also that $\tilde{\omega} := \frac{i}{2}(\varphi^{1\bar{1}} + \varphi^{2\bar{2}} + \varphi^{3\bar{3}})$ is not d-closed but $d\tilde{\omega}^2 = 0$, from which it follows that the manifold M^6 admits a balanced metric.

Moreover, since M^6 is a compact quotient of a completely-solvable Lie group, by the K. Hasegawa's theorem [18, Main Theorem], we have the following result, see also [14, Theorem 3.3]. (We recall that a compact complex manifold is said to belong to *class C of Fujiki* if it admits a proper modification from a Kähler manifold.)

Theorem 3.2 ([18, Main Theorem]). *The manifold M^6 admits no Kähler structure and it is not in class C of Fujiki.*

3.3. An almost-Kähler structure on the Nakamura manifold. By K. Hasegawa's theorem [18, Main Theorem], any integrable complex structure on M^6 (for example, the J defined in §3.2) does not admit any symplectic structure compatible with it. Therefore, we consider the almost-complex structure J' defined by

$$J'e^1 := -e^2, \quad J'e^3 := -e^4, \quad J'e^5 := -e^6;$$

considering

$$\psi^1 := \frac{1}{2}(e^1 + ie^2), \quad \psi^2 := e^3 + ie^4, \quad \psi^3 := e^5 + ie^6$$

as a co-frame for the space of $(1,0)$ -forms on (M^6, J') , one can compute

$$d\psi^1 = 0, \quad d\psi^2 = \psi^{1\bar{2}} + \psi^{\bar{1}2}, \quad d\psi^3 = \psi^{1\bar{3}} + \psi^{\bar{1}3},$$

from which it is clear that J' is not integrable. Note that the J' -compatible 2-form

$$\omega' := e^{12} + e^{34} + e^{56}$$

is d-closed. Hence, (M^6, J', ω') is an almost-Kähler manifold.

Moreover, recall that

$$H_{dR}^2(M^6; \mathbb{R}) \simeq \underbrace{\mathbb{R} \langle i\psi^{1\bar{1}}, i\psi^{2\bar{2}}, i\psi^{3\bar{3}}, i(\psi^{2\bar{3}} + \psi^{3\bar{2}}) \rangle}_{\subseteq H_{J'}^+(M^6)_{\mathbb{R}}} \oplus \underbrace{\mathbb{R} \langle i(\psi^{23} - \psi^{\bar{2}\bar{3}}) \rangle}_{\subseteq H_{J'}^-(M^6)_{\mathbb{R}}},$$

where we have listed the harmonic representatives with respect to the metric $g' := \sum_{j=1}^6 e^j \odot e^j$ instead of their classes; note that the listed g' -harmonic representatives are of pure type with respect to J' . Therefore, J' is obviously \mathcal{C}^∞ -full; it is also \mathcal{C}^∞ -pure by [15, Proposition 3.2] or [11, Proposition 2.8], see Proposition 2.1. Moreover, since any cohomology class in $H_{J'}^+(M^6)_\mathbb{R}$ (respectively, in $H_{J'}^-(M^6)_\mathbb{R}$) has a g' -harmonic representative in $\mathcal{Z}_{J'}^{(1,1)}$ (respectively, in $\mathcal{Z}_{J'}^{(2,0),(0,2)}$), by [15, Theorem 3.7] we have that J' is also pure-and-full. One can explicitly check that the Lefschetz type operator $\mathcal{L}_{\omega'}: \wedge^2 M^6 \rightarrow \wedge^4 M^6$ introduced in §2 takes g' -harmonic 2-forms to g' -harmonic 4-forms, since

$$\begin{aligned} \mathcal{L}_{\omega'} e^{12} &= e^{1234} + e^{1256} = *_{g'} (e^{34} + e^{56}) , & \mathcal{L}_{\omega'} e^{36} &= e^{1236} = *_{g'} e^{45} , \\ \mathcal{L}_{\omega'} e^{34} &= e^{1234} + e^{3456} = *_{g'} (e^{12} + e^{56}) , & \mathcal{L}_{\omega'} e^{45} &= e^{1245} = *_{g'} e^{36} , \\ \mathcal{L}_{\omega'} e^{56} &= e^{1256} + e^{3456} = *_{g'} (e^{12} + e^{34}) . \end{aligned}$$

Resuming, we have shown the following result.

Proposition 3.3. *Let M^6 be the Nakamura manifold. Then there exist a complex structure J and an almost-Kähler structure (J', ω', g') , both of which are \mathcal{C}^∞ -pure-and-full and pure-and-full.*

Furthermore, the Lefschetz type operator of the almost-Kähler structure (J', ω', g') takes g' -harmonic 2-forms to g' -harmonic 4-forms.

Inspired by the argument of the proof of [11, Theorem 2.3], see Theorem 1.3, one can ask the following question, compare also [13, §2]; we provide in Proposition 4.1 an example of a non- \mathcal{C}^∞ -full almost-Kähler structure for which the Lefschetz type property on 2-forms does not hold.

Question 3.4. Let (M, J, ω, g) be a compact $2n$ -dimensional almost-Kähler manifold satisfying the Lefschetz type property on 2-forms. Is J \mathcal{C}^∞ -full?

4. AN ALMOST-KÄHLER NON- \mathcal{C}^∞ -FULL STRUCTURE

Let $X := \mathbb{Z}[i]^3 \setminus (\mathbb{C}^3, *)$ be the *Iwasawa manifold*, where the group structure on \mathbb{C}^3 is defined by

$$(z_1, z_2, z_3) * (w_1, w_2, w_3) := (z_1 + w_1, z_2 + w_2, z_3 + z_1 w_2 + w_3) .$$

Considering the standard complex structure induced by the one on \mathbb{C}^3 and setting $\{\varphi^1, \varphi^2, \varphi^3\}$ as a global co-frame for the $(1, 0)$ -forms on X , by A. Hattori's theorem [19, Corollary 4.2], see Theorem 5.3, one gets that

$$\begin{aligned} H_{dR}^2(X; \mathbb{C}) &\simeq \mathbb{R} \left\langle \varphi^{13} + \varphi^{\bar{1}\bar{3}}, i(\varphi^{13} - \varphi^{\bar{1}\bar{3}}), \varphi^{23} + \varphi^{\bar{2}\bar{3}}, i(\varphi^{23} - \varphi^{\bar{2}\bar{3}}), \right. \\ &\quad \left. \varphi^{12} - \varphi^{2\bar{1}}, i(\varphi^{12} + \varphi^{2\bar{1}}), i\varphi^{1\bar{1}}, i\varphi^{2\bar{2}} \right\rangle \otimes_{\mathbb{R}} \mathbb{C} , \end{aligned}$$

where we have listed the harmonic representatives with respect to the metric $g := \sum_{h=1}^3 \varphi^h \odot \bar{\varphi}^h$ instead of their classes. Set

$$\varphi^1 =: e^1 + i e^2, \quad \varphi^2 =: e^3 + i e^4, \quad \varphi^3 =: e^5 + i e^6;$$

then,

$$d e^5 = -e^{13} + e^{24}, \quad d e^6 = -e^{14} - e^{23},$$

the other differentials being zero. Therefore,

$$H_{dR}^2(X; \mathbb{R}) \simeq \mathbb{R} \langle e^{15} - e^{26}, e^{16} + e^{25}, e^{35} - e^{46}, e^{36} + e^{45}, e^{13} + e^{24}, e^{23} - e^{14}, e^{12}, e^{34} \rangle .$$

Set

$$\begin{aligned} v_1 &:= e^{15} - e^{26}, & v_2 &:= e^{16} + e^{25}, & v_3 &:= e^{35} - e^{46}, & v_4 &:= e^{36} + e^{45}, \\ v_5 &:= e^{13} + e^{24}, & v_6 &:= e^{23} - e^{14}, & v_7 &:= e^{12}, & v_8 &:= e^{34}. \end{aligned}$$

Consider the almost-Kähler structure (J, ω, g) on X defined by

$$Je^1 := -e^6, \quad Je^2 := -e^5, \quad Je^3 := -e^4, \quad \omega := e^{16} + e^{25} + e^{34}.$$

We easily get that

$$\mathbb{R} \langle v_2, v_3 + v_5, v_4 - v_6, v_8 \rangle \subseteq H_J^+(X), \quad \mathbb{R} \langle v_1, v_3 - v_5, v_4 + v_6 \rangle \subseteq H_J^-(X).$$

We claim that the previous inclusions are actually equalities, and in particular that J is a non- \mathcal{C}^∞ -full almost-Kähler structure on X .

Indeed, we firstly note that, by [15, Proposition 3.2] or [11, Proposition 2.8], see Proposition 2.1, J is \mathcal{C}^∞ -pure, since it admits a symplectic structure compatible with it. Moreover, we recall that a \mathcal{C}^∞ -full almost-complex structure is also pure by [20, Proposition 2.30] and therefore it satisfies also that

$$(1) \quad H_J^{(3,1),(1,3)}(X)_{\mathbb{R}} \cap H_J^{(2,2)}(X)_{\mathbb{R}} = \{0\},$$

see [1, Theorem 2.4]. Therefore, our claim reduces to prove that J does not satisfy (1). Note that

$$\begin{aligned} [e^{3456}] &= [e^{3456} - d e^{135}] = [e^{3456} + e^{1234}] \\ &= [e^{3456} + d e^{135}] = [e^{3456} - e^{1234}] \end{aligned}$$

and that $e^{3456} + e^{1234} \in \wedge_J^{(3,1),(1,3)}(X)_{\mathbb{R}}$ while $e^{3456} - e^{1234} \in \wedge_J^{(2,2)}(X)_{\mathbb{R}}$, and so $H_J^{(3,1),(1,3)}(X)_{\mathbb{R}} \cap H_J^{(2,2)}(X)_{\mathbb{R}} \ni [e^{3456}]$, therefore (1) does not hold, and hence J is not \mathcal{C}^∞ -full.

Let \mathcal{L}_ω be the Lefschetz type operator of the almost-Kähler structure (J, ω, g) . Then, we have $\mathcal{L}_\omega(e^{12}) = e^{1234} = d(e^{245})$, i.e., \mathcal{L}_ω does not take g -harmonic 2-forms in g -harmonic 4-forms.

Hence, we have proved the following result.

Proposition 4.1. *Let $X := \mathbb{Z}[\mathfrak{i}]^3 \setminus (\mathbb{C}^3, *)$ be the real manifold underlying the Iwasawa manifold. Then there exists an almost-Kähler structure (J, ω, g) on X which is \mathcal{C}^∞ -pure and non- \mathcal{C}^∞ -full.*

Furthermore, the Lefschetz type operator of the almost-Kähler structure (J, ω, g) does not take g -harmonic 2-forms to g -harmonic 4-forms.

5. ALMOST-COMPLEX MANIFOLDS WITH LARGE ANTI-INVARIANT COHOMOLOGY

Given an almost-complex structure J on a compact manifold M , it is natural to ask how large the cohomology subgroup $H_J^{(2,0),(0,2)}(M)_{\mathbb{R}}$ can be. In this direction, T. Drăghici, T.-J. Li, and the third author raised the following question in [12].

Question 5.1 ([12, Conjecture 2.5]). *Are there compact 4-dimensional manifold M endowed with non-integrable almost-complex structures J such that $\dim_{\mathbb{R}} H_J^-(M) \geq 3$?*

We present here a 1-parameter family $\{J_t\}_t$ of (non-integrable) almost-complex structures on the 6-dimensional torus \mathbb{T}^6 having $h_{J_t}^- := \dim_{\mathbb{R}} H_{J_t}^-(\mathbb{T}^6)_{\mathbb{R}}$ greater than 3, see also [1, §4]. For t small enough, set $\alpha_t := \alpha_t(x^3) \in \mathcal{C}^\infty(\mathbb{T}^6)$ such that $\alpha_0(x^3) \equiv 1$ and set

$$\varphi_t^1 := dx^1 + i \alpha_t dx^4, \quad \varphi_t^2 := dx^2 + i dx^5, \quad \varphi_t^3 := dx^3 + i dx^6;$$

therefore, the structure equations are

$$d\varphi_t^1 = i d\alpha_t \wedge dx^4, \quad d\varphi_t^2 = 0, \quad d\varphi_t^3 = 0.$$

Straightforward computations give that the J -anti-invariant d-closed 2-forms are of the type

$$\psi = \frac{C}{\alpha_t} (dx^{13} - \alpha_t dx^{46}) + D (dx^{16} - \alpha_t dx^{34}) + E (dx^{23} - dx^{56}) + F (dx^{26} - dx^{35}),$$

where $C, D, E, F \in \mathbb{R}$ (we shorten $dx^j \wedge dx^k$ by dx^{jk}). Moreover, the forms $dx^{23} - dx^{56}$ and $dx^{26} - dx^{35}$ are clearly harmonic with respect to the standard flat metric $\sum_{j=1}^6 dx^j \otimes dx^j$, while the classes of $dx^{16} - \alpha_t dx^{34}$ and $dx^{13} - \alpha_t dx^{46}$ are non-zero, their harmonic parts being non-zero. Hence, we get that $h_{J_0}^- = 6$ and

$$h_{J_t}^- = 4 \quad \text{for small } t \neq 0.$$

In the general case, we ask the following natural question.

Question 5.2. Are there examples of non-integrable almost-complex structures J on a compact $2n$ -dimensional manifold with $\dim_{\mathbb{R}} H_J^-(M) > n(n-1)$?

Consider now a solvmanifold $M = \Gamma \backslash G$, namely, a compact quotient of a connected simply-connected solvable Lie group G by a co-compact discrete subgroup Γ . Denote the Lie algebra naturally associated to G by \mathfrak{g} , and consider $(\wedge^\bullet \mathfrak{g}^*, d)$ the subcomplex of the de Rham complex $(\wedge^\bullet M, d)$ given by the left-invariant differential forms. The following result by K. Nomizu [25] and A. Hattori [19] holds.

Theorem 5.3 ([25, Theorem 1], [19, Theorem 4.2]). *Let M be a nilmanifold or, more in general, a completely-solvable solvmanifold. Then $H^\bullet(\wedge^\bullet \mathfrak{g}^*, d) \simeq H_{dR}^\bullet(M; \mathbb{R})$.*

Let J be a left-invariant almost-complex structure on M , namely, an almost-complex structure on M induced by an almost-complex structure on G that is invariant under the action of G on itself given by left-translations. Given $p, q \in \mathbb{N}$, denote by

$$H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}} := \left\{ \mathfrak{a} = [\alpha] \in H^\bullet(\wedge^\bullet \mathfrak{g}^*, d) : \alpha \in \wedge_J^{(p,q),(q,p)} \mathfrak{g}^* \right\} \subseteq H_{dR}^\bullet(M; \mathbb{R})$$

the subgroup (see, e.g., [8, Lemma 9]) of $H_{dR}^\bullet(M; \mathbb{R})$ that consists of classes admitting a left-invariant representative of type $(p, q) + (q, p)$, where $\wedge_J^{(p,q),(q,p)} \mathfrak{g}^* := (\wedge^{p,q}(\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C})^* \oplus \wedge^{q,p}(\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C})^*) \cap \wedge^\bullet \mathfrak{g}^*$.

Using Belgun's symmetrization trick, [4, Theorem 7], one can prove the following Nomizu-type result, which relates the subgroups $H_J^{(p,q),(q,p)}(M)_{\mathbb{R}}$ with their left-invariant part $H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}}$.

Theorem 5.4. *Let $M = \Gamma \backslash G$ be a solvmanifold endowed with a left-invariant almost-complex structure J , and denote the Lie algebra naturally associated to G by \mathfrak{g} . For any $p, q \in \mathbb{N}$, the map*

$$j: H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}} \rightarrow H_J^{(p,q),(q,p)}(M)_{\mathbb{R}}$$

induced by left-translations is injective, and, if $H_{dR}^{\bullet}(\wedge^{\bullet} \mathfrak{g}^, d) \simeq H_{dR}^{\bullet}(M; \mathbb{R})$ (for instance, if M is a completely-solvable solvmanifold), then $j: H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}} \rightarrow H_J^{(p,q),(q,p)}(M)_{\mathbb{R}}$ is in fact an isomorphism.*

Proof. Since J is left-invariant, left-translations induce the map $j: H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}} \rightarrow H_J^{(p,q),(q,p)}(M)_{\mathbb{R}}$.

Since, by J. Milnor's Lemma [24, Lemma 6.2], G is unimodular, one can take in particular a bi-invariant volume form η on M such that $\int_M \eta = 1$. Consider the F. A. Belgun symmetrization map in [4, Theorem 7], namely,

$$\mu: \wedge^{\bullet} M \rightarrow \wedge^{\bullet} \mathfrak{g}^*, \quad \mu(\alpha) := \int_M \alpha|_m \eta(m).$$

Since μ commutes with d by [4, Theorem 7], it induces the map $\mu: H_{dR}^{\bullet}(M; \mathbb{R}) \rightarrow H^{\bullet}(\wedge^{\bullet} \mathfrak{g}^*, d)$, and, since μ commutes with J , it preserves the bi-graduation; therefore it induces the map $\mu: H_J^{(p,q),(q,p)}(M) \rightarrow H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}}$. Moreover, since μ is the identity on the space of left-invariant forms by [4, Theorem 7], we get the commutative diagram

$$\begin{array}{ccccc} H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}} & \xrightarrow{j} & H_J^{(p,q),(q,p)}(M)_{\mathbb{R}} & \xrightarrow{\mu} & H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}} \\ & \searrow & \text{id} & \nearrow & \\ & & & & \end{array}$$

hence $j: H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}} \rightarrow H_J^{(p,q),(q,p)}(M)_{\mathbb{R}}$ is injective, and $\mu: H_J^{(p,q),(q,p)}(M)_{\mathbb{R}} \rightarrow H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}}$ is surjective.

Furthermore, when $H^{\bullet}(\wedge^{\bullet} \mathfrak{g}^*, d) \simeq H_{dR}^{\bullet}(M; \mathbb{R})$ (for instance, when M is a completely-solvable solvmanifold, by A. Hattori's theorem [19, Theorem 4.2], see Theorem 5.3), since $\mu|_{\wedge^{\bullet} \mathfrak{g}^*} = \text{id}|_{\wedge^{\bullet} \mathfrak{g}^*}$ by [4, Theorem 7], we get that $\mu: H_{dR}^{\bullet}(M; \mathbb{R}) \rightarrow H^{\bullet}(\wedge^{\bullet} \mathfrak{g}^*, d)$ is the identity map, and hence $\mu: H_J^{(p,q),(q,p)}(M)_{\mathbb{R}} \rightarrow H_J^{(p,q),(q,p)}(\mathfrak{g})_{\mathbb{R}}$ is also injective, and hence an isomorphism. \square

In particular, if $M = \Gamma \backslash G$ is a $2n$ -dimensional completely-solvable solvmanifold endowed with a left-invariant almost-complex structure J , then

$$\dim_{\mathbb{R}} H_J^{-}(M) \leq n(n-1) \quad \text{and} \quad \dim_{\mathbb{R}} H_J^{+}(M) \leq n^2;$$

this provides a partial negative answer to Question 5.2.

REFERENCES

- [1] D. Angella, A. Tomassini, On cohomological decomposition of almost-complex manifolds and deformations, *J. Symplectic Geom.* **9** (2011), no. 3, 403–428.
- [2] D. Angella, A. Tomassini, On the cohomology of almost-complex manifolds, *Internat. J. Math.* **23** (2012), no. 2, 1250019, 25 pp.
- [3] D. Angella, A. Tomassini, On the $\partial\bar{\partial}$ -Lemma and Bott-Chern cohomology, to appear in *Invent. Math.*, DOI: 10.1007/s00222-012-0406-3.
- [4] F. A. Belgun, On the metric structure of non-Kähler complex surfaces, *Math. Ann.* **317** (2000), no. 1, 1–40.

- [5] Ch. Benson, C. S. Gordon, Kähler and symplectic structures on nilmanifolds, *Topology* **27** (1988), no. 4, 513–518.
- [6] J.-L. Brylinski, A differential complex for Poisson manifolds, *J. Differential Geom.* **28** (1988), no. 1, 93–114.
- [7] G. R. Cavalcanti, New aspects of the dd^c -lemma, Oxford University D. Phil thesis, [arXiv:math/0501406v1](https://arxiv.org/abs/math/0501406v1) [math.DG].
- [8] S. Console, A. Fino, Dolbeault cohomology of compact nilmanifolds, *Transform. Groups* **6** (2001), no. 2, 111–124.
- [9] P. Deligne, Ph. Griffiths, J. Morgan, D. Sullivan, Real homotopy theory of Kähler manifolds, *Invent. Math.* **29** (1975), no. 3, 245–274.
- [10] S. K. Donaldson, *Two-forms on four-manifolds and elliptic equations*, Inspired by S. S. Chern, 153–172, Nankai Tracts Math., **11**, World Sci. Publ., Hackensack, NJ, 2006.
- [11] T. Drăghici, T.-J. Li, W. Zhang, Symplectic forms and cohomology decomposition of almost complex four-manifolds, *Int. Math. Res. Not. IMRN* **2010** (2010), no. 1, 1–17.
- [12] T. Drăghici, T.-J. Li, W. Zhang, On the J -anti-invariant cohomology of almost complex 4-manifolds, *Quart. J. Math.* **00** (2011), DOI: 10.1093/qmath/har0341, 1–29.
- [13] T. Drăghici, T.-J. Li, W. Zhang, Geometry of tamed almost complex structures in dimension 4, ICCM2010 Proceedings, AMS/IP Vol. **51**, 2012, 233–251.
- [14] M. Fernández, V. Muñoz, J. A. Santisteban, Cohomologically Kähler manifolds with no Kähler metrics, *Int. J. Math. Sci.* **2003** (2003), no. 52, 3315–3325.
- [15] A. Fino, A. Tomassini, On some cohomological properties of almost complex manifolds, *J. of Geom. Anal.* **20** (2010), no. 1, 107–131.
- [16] M. Gualtieri, Generalized complex geometry, *Ann. of Math. (2)* **174** (2011), no. 1, 75–123.
- [17] K. Hasegawa, Minimal models of nilmanifolds, *Proc. Am. Math. Soc.* **106** (1989), no. 1, 65–71.
- [18] K. Hasegawa, A note on compact solvmanifolds with Kähler structures, *Osaka J. Math.* **43** (2006), no. 1, 131–135.
- [19] A. Hattori, Spectral sequence in the de Rham cohomology of fibre bundles, *J. Fac. Sci. Univ. Tokyo Sect. I* **8** (1960), no. 1960, 289–331.
- [20] T.-J. Li, W. Zhang, Comparing tamed and compatible symplectic cones and cohomological properties of almost complex manifolds, *Comm. Anal. Geom.* **17** (2009), no. 4, 651–683.
- [21] A. I. Mal'tsev, On a class of homogeneous spaces, *Izvestiya Akad. Nauk. SSSR. Ser. Mat.* **13** (1949), no. 1, 9–32, English translation in *Amer. Math. Soc. Translation: Series 1* **1951** (1951), no. 39, 193–206.
- [22] O. Mathieu, Harmonic cohomology classes of symplectic manifolds, *Comment. Math. Helv.* **70** (1995), no. 1, 1–9.
- [23] S. A. Merkulov, Formality of canonical symplectic complexes and Frobenius manifolds, *Int. Math. Res. Not.* **1998** (1998), no. 14, 727–733.
- [24] J. Milnor, Curvature of left-invariant metrics on Lie groups, *Adv. Math.* **21** (1976), no. 3, 293–329.
- [25] K. Nomizu, On the cohomology of compact homogeneous spaces of nilpotent Lie groups, *Ann. of Math. (2)* **59** (1954), no. 3, 531–538.
- [26] J. Streets, G. Tian, A parabolic flow of pluriclosed metrics, *Int. Math. Res. Not. IMRN* **2010** (2010), no. 16, 3101–3133.
- [27] D. Yan, Hodge structure on symplectic manifolds, *Adv. Math.* **120** (1996), no. 1, 143–154.

(Daniele Angella) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, 56127, PISA, ITALY

E-mail address: angella@mail.dm.unipi.it

(Adriano Tomassini) DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI PARMA, PARCO AREA DELLE SCIENZE 53/A, 43124, PARMA, ITALY

E-mail address: adriano.tomassini@unipr.it

(Weiyi Zhang) UNIVERSITY OF MICHIGAN, DEPARTMENT OF MATHEMATICS, 1825 EAST HALL, ANN ARBOR, MI 48109

E-mail address: wyzhang@umich.edu